

PLANE CURVES AS PFAFFIANS

ANITA BUCKLEY, TOMAŽ KOŠIR

ABSTRACT. Let C be a smooth curve in $\mathbb{P}^2$ given by an equation $F = 0$ of degree d . In this paper we parametrise all linear pfaffian representations of F by an open subset in the moduli space $M_C(2, K_C)$. We construct an explicit correspondence between pfaffian representations of C and rank 2 vector bundles on C with canonical determinant and no sections.

1. INTRODUCTION

Let k be an algebraically closed field and C an irreducible curve defined by a polynomial F of degree d in $\mathbb{P}^2$. When C is smooth its genus equals

$$g = \frac{1}{2}(d-1)(d-2).$$

These notes consider the following question. For a given homogeneous polynomial $F(x_0, x_1, x_2)$ of degree d find a $2d \times 2d$ skew-symmetric matrix

$$A = \begin{bmatrix} 0 & L_{12} & L_{13} & \cdots & L_{12d} \\ -L_{12} & 0 & L_{23} & \cdots & L_{22d} \\ -L_{13} & -L_{23} & 0 & & \\ \vdots & \vdots & & \ddots & \vdots \\ -L_{12d} & -L_{22d} & & \cdots & 0 \end{bmatrix}$$

with linear forms $L_{ij} = a_{ij}^0 x_0 + a_{ij}^1 x_1 + a_{ij}^2 x_2$ such that

$$\text{Pf } A(x_0, x_1, x_2) = cF(x_0, x_1, x_2) \quad \text{for some } c \in k, c \neq 0.$$

Matrix A is called *linear pfaffian representation* of C . Its cokernel is a rank 2 vector bundle on C . Throughout the paper we equate the notion of vector bundles and locally free sheaves.

Two pfaffian representations A and A' are *equivalent* if there exists $X \in \text{GL}_{2d}(k)$ such that

$$A' = XAX^t.$$

A locally free sheaf $\mathcal{E}$ of rank 2 is *stable* if for every invertible sheaf $\mathcal{F} \rightarrow \mathcal{F} \rightarrow 0$ holds

$$\deg \mathcal{F} > \frac{1}{2} \deg \mathcal{E}.$$

Replacing $>$ by $\geq$ defines *semistable*.

We find **all** linear pfaffian representations of C (up to equivalence) and relate them to the moduli space $M_C(2, K_C)$ of semistable rank 2 vector bundles on C with canonical determinant. An explicit construction of representations from the global sections of rank 2 vector bundles with certain properties is given.

In general the elements of A can be homogenous polynomials of various degrees. Such pfaffian representations are considered in Theorem 3.7. A good survey of the linear algebra of Pfaffians can be found in [6, Appendix D]. In [7, Chapter V. 2]

Hartshorne identifies the theory of rank 2 sheaves with the theory of ruled surfaces over C .

Study of pfaffian representations is strongly related to and motivated by determinantal representations. A *linear determinantal representation* of C is a $d \times d$ matrix of linear forms

$$M = x_0 M_0 + x_1 M_1 + x_2 M_2$$

satisfying

$$\det M = cF,$$

where $M_0, M_1, M_2 \in \text{Mat}_d(k)$ and $c \in k, c \neq 0$. Two determinantal representations M and M' are *equivalent* if there exist $X, Y \in \text{GL}_d(k)$ such that

$$M' = XMY.$$

There are many more pfaffian than determinantal representations. Indeed, every determinantal representation M induces *decomposable pfaffian representation*

$$\begin{bmatrix} 0 & M \\ -M^t & 0 \end{bmatrix}.$$

Note that the equivalence relation is well defined since

$$\begin{bmatrix} 0 & XMY \\ -(XMY)^t & 0 \end{bmatrix} = \begin{bmatrix} X & 0 \\ 0 & Y^t \end{bmatrix} \begin{bmatrix} 0 & M \\ -M^t & 0 \end{bmatrix} \begin{bmatrix} X^t & 0 \\ 0 & Y \end{bmatrix}.$$

A brief outline of the paper is the following. In Section 2 we recall the parametrisation of linear determinantal representations by points on the Jacobian variety due to Vinnikov [18]. We use similar ideas in Sections 3 and 4 to parametrise all linear pfaffian representations by points in an open subset of the moduli space $M_C(2, K_C)$. In the third section we give an explicit construction of the correspondence between pfaffian representations and vector bundles with certain properties. In the fourth section we relate these vector bundles to the points on $M_C(2, K_C)$ not on the subvariety cut out by Cartier divisor Θ_{2, K_C} . Section 5 considers decomposable bundles which are parametrised by an open set in Kummer variety. Pfaffian representation arising from a decomposable bundle $\mathcal{F} \oplus (\mathcal{F}^{-1} \otimes \mathcal{O}_C(d-1))$ is exactly the decomposable representation

$$\begin{bmatrix} 0 & M \\ -M^t & 0 \end{bmatrix},$$

where M is the determinantal representation corresponding to $\mathcal{F} \cong \text{Coker } M$. Elliptic curves in Section 6 only allow decomposable representations. In the last section we compute pfaffian representations of a genus 3 curve. To our knowledge there are not many results explicitly describing $M_C(2, K_C)$ for curves of higher genus. Methods considered in this paper could be helpful in finding all pfaffian representations and consequently the moduli space of $g > 3$ curves.

2. DETERMINANTAL REPRESENTATIONS AND THE JACOBIAN

There is a one to one correspondence between linear determinantal representations (up to equivalence) of C and line bundles (up to isomorphism) on C with certain properties. This well known result is summed up in the following theorem of Beauville [1, Proposition 3.1].

Theorem 2.1. *Let C be a plane curve defined by a polynomial F of degree d and let L be a line bundle of degree $\frac{1}{2}d(d-1)$ on C with $H^0(C, L(-1)) = 0$. Then there exists a $d \times d$ linear matrix M with $\det M = F$ and an exact sequence*

$$(1) \quad 0 \rightarrow \bigoplus_{i=1}^d \mathcal{O}_{\mathbb{P}^2}(-1) \xrightarrow{M} \bigoplus_{i=1}^d \mathcal{O}_{\mathbb{P}^2} \rightarrow L \rightarrow 0.$$

Conversely, let M be a linear $d \times d$ matrix with $\det M = F$. Then its cokernel is a line bundle of degree $\frac{1}{2}d(d-1)$ and $H^0(C, \text{Coker } M(-1)) = 0$.

Dolgachev explicitly described the above correspondence in [4]. Vinnikov also gave an explicit construction of the correspondence in [18, Theorems 2-4]. Additionally he related determinantal representations to points on the Jacobian variety in the following way.

It is known [3, Theorem 1.1] that representations M and M' are equivalent if and only if $\text{Coker } M$ and $\text{Coker } M'$ are isomorphic sheaves. Therefore the problem of classifying all linear representations of F (up to equivalence) is the same as finding all line bundles with the property $\deg L = \frac{1}{2}d(d-1)$ and $H^0(C, L(-1)) = 0$. In order to simplify the notation we tensor the above by $\mathcal{O}(1)$ and consider line bundles with $\deg \mathcal{L} = \frac{1}{2}d(d-1) - d = \frac{1}{2}d(d-3)$ and $H^0(C, \mathcal{L}) = 0$. We call the bundles with this property *nonexceptional* line bundles. Analogously, line bundles of degree $\frac{1}{2}d(d-3)$ and $H^0(C, \mathcal{L}) \neq 0$ are called *exceptional*.

Lemma 2.2. *Let C be of genus g . The exceptional divisor classes define a $g-1$ dimensional subvariety W_{g-1} on the Jacobian variety J .*

This proves the following theorem.

Theorem 2.3. *All linear determinantal representations of F (up to equivalence) can be parametrised by points on the Jacobian variety of C not on the exceptional subvariety W_{g-1} .*

3. CLASSIFICATION OF PFAFFIAN REPRESENTATIONS FROM THE SCRATCH

In this section we consider the following problem. For a given homogeneous polynomial $F(x_0, x_1, x_2)$ of degree d find all linear pfaffian representations. The main result of this section is an elementary proof of Theorem 3.6 and explicit construction of representations from suitable vector bundles.

Following the ideas of Dolgachev [4] formulate the problem geometrically and coordinate free. Let U be a $2d$ dimensional vector space. It is well known that $\bigwedge^2 U$ can be identified with $2d \times 2d$ skew-symmetric matrices. Let Ω_k denote the set of vectors $\sum_{i=1}^k v_i \wedge w_i$ in $\bigwedge^2 U$ where $\dim\{v_1, \dots, v_k, w_1, \dots, w_k\} = 2k$. Elements of Ω_k are said to have *irreducible length k* since they can be written as a sum of k and not less than k pure nonzero products in $\bigwedge^2 U$. In [20] it is shown that

Lemma 3.1. *Ω_k is isomorphic to the set of all rank $2k$ skew-symmetric matrices.*

Proof. Specifically, let $e_1, \dots, e_{2d}$ be a basis for U and $\{E_{ij}\}$ the standard basis for $2d \times 2d$ matrices. Then the isomorphism equals

$$e_i \wedge e_j \mapsto E_{ij} - E_{ji}$$

and extends linearly to

$$(2) \quad \sum_{i=1}^{2d} \alpha_i e_i \wedge \sum_{j=1}^{2d} \beta_j e_j \mapsto \sum_{i,j=1}^{2d} (\alpha_i \beta_j - \alpha_j \beta_i) (E_{ij} - E_{ji}).$$

□

Next let E be a 3 dimensional vector space and Φ a linear map

$$\Phi: \mathbb{P}(E) \longrightarrow \mathbb{P}(\bigwedge^2 U).$$

Note that Φ corresponds to a skew-symmetric matrix with linear forms as its elements. Alternatively, Φ is an element in $E^* \otimes (\bigwedge^2 U)$.

Let $\mathcal{P}_d \subset \mathbb{P}(\bigwedge^2 U)$ be the hypersurface parametrising non-invertible linear maps. Choose a basis of U , then $\mathcal{P}_d$ is given by the pfaffian of a skew-symmetric matrix. The inverse image of $\mathcal{P}_d$ under Φ is a plane curve of degree d in $\mathbb{P}(E)$.

Let C be a smooth plane curve defined by F . Assume that C admits a pfaffian representation A .

Lemma 3.2. *For any $x \in C$ the corank of $A(x)$ equals 2.*

Proof. Assume

$$A = [a_{ij}^0 x_0 + a_{ij}^1 x_1 + a_{ij}^2 x_2] \quad \text{and} \quad \text{Pf } A = cF, \quad c \neq 0.$$

Denote by $\text{Pf}^{ij} A$ the pfaffian of the $(2d-2) \times (2d-2)$ skew-symmetric matrix obtained by removing the i th and j th rows and columns from A . Then

$$\frac{\partial F}{\partial x_k}(x) = \frac{1}{c} \sum_{i,j} a_{ij}^k \text{Pf}^{ij} A(x).$$

If for some $x \in C$ all $2d-2$ pfaffian minors vanish, then x must be a singular point of F . By our assumption F is smooth, thus $\text{rank } A(x) \geq 2d-2$ for all $x \in C$. This ends the proof because rank of skew-symmetric matrices is even and $\det A = F^2 = 0$. □

Define the *pfaffian adjoint* of A to be the skew-symmetric matrix

$$\tilde{A} = \begin{bmatrix} 0 & & & \\ & \ddots & & \\ & & (-1)^{i+j} \text{Pf}^{ij} A & \\ & & & \ddots \\ & & & & 0 \end{bmatrix}.$$

By analogy with determinants the following holds

$$(3) \quad \tilde{A} \cdot A = \text{Pf } A \cdot \text{Id}_{2d}.$$

More properties and linear algebra of Pfaffians can be found in [6, Appendix D].

By Lemma 3.2 the cokernel of A defines a rank 2 bundle over C . Since $\text{Pf } A = cF$ the cokernel can be obtained from $\tilde{A}$ by (3). The properties of $\text{Coker } A$ are described in the following proposition.

Proposition 3.3. *Let A be a pfaffian representation of a smooth plane curve C defined by a homogeneous polynomial F of degree d . Then $\mathcal{E} = \text{Coker } A$ is a rank 2 vector bundle on C and*

- (i) $h^0(C, \mathcal{E}) = 2d$,
- (ii) $H^0(C, \mathcal{E}(-1)) = H^1(C, \mathcal{E}(-1)) = 0$,

$$(iii) \det \mathcal{E} = \bigwedge^2 \mathcal{E} = \mathcal{O}_C(d-1).$$

Proof. By definition $\mathcal{E}$ fits into the exact sequence

$$(4) \quad 0 \rightarrow \bigoplus_{i=1}^{2d} \mathcal{O}_{\mathbb{P}^2}(-1) \xrightarrow{A} \bigoplus_{i=1}^{2d} \mathcal{O}_{\mathbb{P}^2} \rightarrow \mathcal{E} \rightarrow 0.$$

Applying the functor $H^i(\mathbb{P}^2, *)$ on (4) gives

$$H^0(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}(-1)^{2d}) \rightarrow H^0(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}^{2d}) \rightarrow H^0(\mathbb{P}^2, \mathcal{E}) \rightarrow H^1(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}(-1)^{2d}) \rightarrow \dots$$

$$\begin{array}{ccc} 0 & 2d & 0 \end{array}$$

where the bottom row denotes dimensions of the cohomology of projective space which is computed in [7, Theorem III.5.1]. Thus $\dim H^0(\mathbb{P}^2, \mathcal{E}) = 2d$.

Next tensor (4) by $\mathcal{O}_{\mathbb{P}^2}(-1)$ and again apply the functor $H^i(\mathbb{P}^2, *)$. This gives a long exact sequence with dimensions

$$\begin{array}{ccccc} H^0(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}(-2)^{2d}) & \rightarrow & H^0(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}(-1)^{2d}) & \rightarrow & H^0(\mathbb{P}^2, \mathcal{E}(-1)) \rightarrow \\ \parallel & & \parallel & & \\ 0 & & 0 & & \\ \\ H^1(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}(-2)^{2d}) & \rightarrow & H^1(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}(-1)^{2d}) & \rightarrow & H^1(\mathbb{P}^2, \mathcal{E}(-1)) \rightarrow \\ \parallel & & \parallel & & \\ 0 & & 0 & & \\ \\ H^2(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}(-2)^{2d}) & \rightarrow & \dots & & \\ \parallel & & & & \\ 0 & & & & \end{array}$$

where $H^2(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}(-2)^{2d}) \cong H^0(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}(-1)^{2d}) = 0$ by Serre duality. We proved that $H^0(\mathbb{P}^2, \mathcal{E}(-1)) = H^1(\mathbb{P}^2, \mathcal{E}(-1)) = 0$.

Since $\mathcal{E}$ is supported on C we proved (i) and (ii).

In order to prove (iii) apply the functor $\mathcal{H}om_{\mathcal{O}_{\mathbb{P}^2}}(*, \mathcal{O}_{\mathbb{P}^2}(-1))$ to (4). We get

$$\begin{aligned} 0 \rightarrow \mathcal{H}om(\mathcal{E}, \mathcal{O}_{\mathbb{P}^2}(-1)) &\rightarrow \mathcal{H}om(\mathcal{O}_{\mathbb{P}^2}^{2d}, \mathcal{O}_{\mathbb{P}^2}(-1)) \rightarrow \mathcal{H}om(\mathcal{O}_{\mathbb{P}^2}(-1)^{2d}, \mathcal{O}_{\mathbb{P}^2}(-1)) \rightarrow \\ &\rightarrow \mathcal{E}xt^1(\mathcal{E}, \mathcal{O}_{\mathbb{P}^2}(-1)) \rightarrow \mathcal{E}xt^1(\mathcal{O}_{\mathbb{P}^2}^{2d}, \mathcal{O}_{\mathbb{P}^2}(-1)) \rightarrow \dots, \end{aligned}$$

where

$$\begin{aligned} \mathcal{H}om(\mathcal{O}_{\mathbb{P}^2}^{2d}, \mathcal{O}_{\mathbb{P}^2}(-1)) &\cong (\mathcal{O}_{\mathbb{P}^2}^{2d})^\vee \otimes \mathcal{O}_{\mathbb{P}^2}(-1) \cong \mathcal{O}_{\mathbb{P}^2}(-1)^{2d}, \\ \mathcal{H}om(\mathcal{O}_{\mathbb{P}^2}(-1)^{2d}, \mathcal{O}_{\mathbb{P}^2}(-1)) &\cong (\mathcal{O}_{\mathbb{P}^2}(-1)^{2d})^\vee \otimes \mathcal{O}_{\mathbb{P}^2}(-1) \cong \mathcal{O}_{\mathbb{P}^2}^{2d} \end{aligned}$$

and $\mathcal{E}xt^1(\mathcal{O}_{\mathbb{P}^2}^{2d}, \mathcal{O}_{\mathbb{P}^2}(-1)) = 0$ by [7, Proposition III.6.3]. This implies that $\mathcal{E}xt^i(\mathcal{E}, \mathcal{O}_{\mathbb{P}^2}(-1)) = 0$ for $i \neq 1$ and the above sequence is isomorphic to

$$(5) \quad 0 \rightarrow \bigoplus_{i=1}^{2d} \mathcal{O}_{\mathbb{P}^2}(-1) \xrightarrow{A^t} \bigoplus_{i=1}^{2d} \mathcal{O}_{\mathbb{P}^2} \rightarrow \mathcal{E}xt^1(\mathcal{E}, \mathcal{O}_{\mathbb{P}^2}(-1)) \rightarrow 0.$$

Thus we obtain

$$\begin{aligned} \mathcal{E} &\cong \text{Coker } A \cong \text{Coker } A^t \cong \mathcal{E}xt^1(\mathcal{E}, \mathcal{O}_{\mathbb{P}^2}(-1)) \cong \\ &\mathcal{E}xt^1(\mathcal{E}, \mathcal{O}_{\mathbb{P}^2}(-3) \otimes \mathcal{O}_{\mathbb{P}^2}(2)) \cong \mathcal{E}xt^1(\mathcal{E}, \mathcal{O}_{\mathbb{P}^2}(-3)) \otimes \mathcal{O}_{\mathbb{P}^2}(2) \cong \\ &\mathcal{H}om_C(\mathcal{E}, \mathcal{O}_C(d-3)) \otimes \mathcal{O}_{\mathbb{P}^2}(2) \cong \mathcal{H}om_C(\mathcal{E}, \mathcal{O}_C(d-1)) \cong \\ &\mathcal{E}^\vee \otimes \mathcal{O}_C(d-1) \end{aligned}$$

since by Serre duality

$$\mathcal{E}xt^1(\mathcal{E}, \mathcal{O}_{\mathbb{P}^2}(-3)) \cong \mathcal{E}xt^1(\mathcal{E}, \omega_{\mathbb{P}^2}) \cong \mathcal{H}om_C(\mathcal{E}, \omega_C) \cong \mathcal{H}om_C(\mathcal{E}, \mathcal{O}_C(d-3)).$$

Finally,

$$\wedge^2 \mathcal{E} \cong \mathcal{E} \wedge (\mathcal{E}^\vee \otimes \mathcal{O}_C(d-1)) \cong \mathcal{O}_C(d-1).$$

□

In the sequel the reverse problem will be considered. We will give an explicit construction of pfaffian representation from a vector bundle with certain properties.

Theorem 3.4. *Let C be a smooth plane curve of degree d . To every rank 2 vector bundle $\mathcal{E}$ on C with properties*

- (i) $h^0(C, \mathcal{E}) = 2d$,
- (ii) $H^0(C, \mathcal{E}(-1)) = 0$,
- (iii) $\det \mathcal{E} = \wedge^2 \mathcal{E} = \mathcal{O}_C(d-1)$

we can assign a pfaffian representation $A_{\mathcal{E}}$. In particular, isomorphic bundles induce equivalent representations.

Proof. Let $U = H^0(C, \mathcal{E})$ be the $2d$ dimensional vector space of global sections of $\mathcal{E}$. We will define a map ψ from C to the space of $2d \times 2d$ skew-symmetric matrices with entries from the space of homogeneous polynomials of degree $d-1$, such that $\psi^{-1}(\mathcal{P}_d) = C$.

Choose a basis $\{s_1, \dots, s_{2d}\}$ for U and define

$$C \ni x \xrightarrow{\psi} \sum_{1 \leq i < j \leq 2d} (s_i(x) \wedge s_j(x))(E_{ij} - E_{ji}) = \begin{bmatrix} 0 & & & \\ & \ddots & s_i(x) \wedge s_j(x) & \\ & & \ddots & \\ & & & 0 \end{bmatrix}.$$

Since $s_i \wedge s_j \in \wedge^2 U$, by property (iii) the map ψ extends to

$$\Psi: \mathbb{P}(E) \longrightarrow \mathbb{P}(\wedge^2 U)$$

given by a linear system of plane curves of degree $d-1$. In other words, Ψ is a tensor in $S^{d-1}(E^*) \otimes (\wedge^2 U)$. In coordinates it equals to a $2d \times 2d$ skew-symmetric matrix $B(x_0, x_1, x_2)$ with entries from the space of homogeneous polynomials of degree $d-1$.

From the definition of B and isomorphism (2) in Lemma 3.1 it follows that at any point $x \in C$ $\text{rank } B(x) = 2$.

Before we proceed, we prove that a different basis $\{s'_1, \dots, s'_{2d}\}$ for U induces equivalent representation B' . Indeed,

$$\begin{bmatrix} s'_1 \\ \vdots \\ s'_{2d} \end{bmatrix} = R \cdot \begin{bmatrix} s_1 \\ \vdots \\ s_{2d} \end{bmatrix}$$

for some constant invertible matrix R , hence

$$B' = \begin{bmatrix} 0 & & & \\ & \ddots & s'_i \wedge s'_j & \\ & & \ddots & \\ & & & 0 \end{bmatrix} = R \cdot \begin{bmatrix} 0 & & & \\ & \ddots & s_i \wedge s_j & \\ & & \ddots & \\ & & & 0 \end{bmatrix} \cdot R^t = R \cdot B \cdot R^t.$$

As before, let F be the defining polynomial for C . Denote by M a 4×4 submatrix of B , obtained by deleting $2d - 4$ rows and columns with the same indexes. Then $F \mid \text{Pf } M$ since $\text{Pf } M(x) = 0$ for all $x \in C$. Consider next a 6×6 skew-symmetric submatrix N of B and its pfaffian adjoint $\tilde{N}$. By (3) we get

$$\begin{aligned} \text{Pf } N \cdot \text{Pf } \tilde{N} &= (\text{Pf } N)^{\text{order } N/2}, \\ \text{Pf } \tilde{N} &= (\text{Pf } N)^{\text{order } N/2-1} = (\text{Pf } N)^2. \end{aligned}$$

The entries of $\tilde{N}$ are Pfaffians of 4×4 submatrices, hence $F^3 \mid (\text{Pf } N)^2$. Since C is irreducible, $F^2 \mid \text{Pf } N$. By repeating this process we obtain that F^{d-2} divides all the Pfaffians of $(2d-2) \times (2d-2)$ skew-symmetric submatrices of B . These are exactly $\text{Pf}^{ij} B$ defined in the proof of Lemma 3.2. This means that

$$A = \frac{1}{F^{d-2}} \tilde{B}$$

is a matrix with entries in E^* . Since $\text{rank } B(x) = 2$ for all $x \in C$, we get $\text{rank } A(x) = 2d - 2$. Therefore $\text{Pf } A$ is a hypersurface of degree d vanishing on C unless $\text{Pf } A$ is identically zero. This implies $\text{Pf } A = cF$ for some constant c .

It remains to prove that $\text{Pf } A \neq 0$. We will use the following remark to describe the sections of $\mathcal{E}(-1)$.

Remark 3.5. Consider the map

$$\begin{aligned} r : C &\longrightarrow \mathbb{P}(U) \\ x &\longmapsto [s_1(x), \dots, s_{2d}(x)]. \end{aligned}$$

Every global section of $\mathcal{E}$ can be seen as a hyperplane in $\mathbb{P}(U)$. A nonzero element in $H^0(C, \mathcal{E}(-1))$ is a section s of $\mathcal{E}$ such that the divisor

$$r(C) \cap (\text{the hyperplane in } \mathbb{P}(U) \text{ defined by } s)$$

equals $r(C \cap L)$ for some line L . All divisors on C which are cut out by a line are equivalent. Hence for any line L the divisor $r(C \cap L)$ on the image $r(C)$ is cut out by a hyperplane in $\mathbb{P}(U)$.

Choose L such that it intersects C in d distinct points $\{p_1, \dots, p_d\}$. Assumption (ii) and Remark 3.5 imply that the divisor $\{r(p_1), \dots, r(p_d)\}$ on $r(C)$ is NOT cut out by a hyperplane in $\mathbb{P}(U)$. This proves that the images

$$\{\Psi(p_1), \dots, \Psi(p_d)\}$$

span a $d - 1$ (projectively) dimensional subspace W in $\mathbb{P}(\bigwedge^2 U)$. Indeed, chose coordinates so that the points $r(p_i) = [s_1(p_i), \dots, s_{2d}(p_i)]$ are defined by

$$\begin{bmatrix} 0 & \dots & 1 & \dots & 0 & 0 & \dots & 0 & \dots & 0 \\ 0 & \dots & 0 & \dots & 0 & 0 & \dots & 1 & \dots & 0 \end{bmatrix}$$

with 1 on the i th and $d + i$ th position. Then W corresponds to the space of $2d \times 2d$ skew-symmetric matrices

$$\begin{bmatrix} 0 & \diamond \\ -\diamond & 0 \end{bmatrix},$$

where $\diamond$ denotes diagonal $d \times d$ matrices.

By (iii) the image $\Psi(L)$ is a Veronese curve of degree $d - 1$. Let l_1, l_2 be the local parameters of L , then $\Psi(L) = l_1^d + *l_1^{d-1}l_2 + \dots + *l_2^d$ is determined by d points. This proves that $\Psi(L)$ lies in W . A general point $\Psi(p) \in \Psi(L)$ does not belong to any hyperplane in W spanned by $d - 1$ points $\Psi(p_i)$. Hence $\Psi(p)$ can be written as a linear combination of the points $\Psi(p_i)$ with nonzero coefficients. In other words,

$\Psi(p)$ can not be written shorter than a sum of d pure nonzero wedge products in $\mathbb{P}(\bigwedge^2 U)$. By Lemma 3.1, $\Psi(p) \in \Omega_d$ corresponds to a skew-symmetric matrix of rank $2d$. Therefore $\text{Pf } B(p) \neq 0$ and hence $\text{Pf } A(p) \neq 0$. $\square$

By the classic result of Cook [3, Theorem 1.1] two representations are equivalent if and only if the corresponding cokernels are isomorphic sheaves. Together with the above considerations it implies

Theorem 3.6. *There is a one to one correspondence between linear pfaffian representations of F (up to equivalence) and rank 2 bundles (up to isomorphism) on C with the property*

$$(6) \quad \det \mathcal{E} \cong \mathcal{O}_C(d-1) \quad \text{and} \quad H^0(C, \mathcal{E}(-1)) = 0.$$

Proof. It remains to show that (i) in Theorem 3.4 follows from conditions (ii) and (iii). For every rank 2 bundle $\mathcal{E} \cong \mathcal{E}^\vee \otimes (\bigwedge^2 \mathcal{E})$ holds. Applying (iii) gives $\mathcal{E} \cong \mathcal{E}^\vee \otimes \mathcal{O}_C(d-1)$. Then

$$\begin{aligned} & H^1(C, \mathcal{E}(-1)) \\ & \cong H^1(C, \mathcal{E}^\vee \otimes \mathcal{O}_C(d-2)) \cong H^0(C, \mathcal{E} \otimes \mathcal{O}_C(2-d) \otimes \mathcal{O}_C(d-3)) \\ & \cong H^0(C, \mathcal{E}(-1)) \end{aligned}$$

by Serre duality. Hence (ii) implies $H^1(X, \mathcal{E}(-1)) = 0$.

Let L be a section of $\mathcal{O}_C(1)$ and consider the exact sequence

$$(7) \quad 0 \longrightarrow \mathcal{E}(-1) \longrightarrow \mathcal{E} \longrightarrow \mathcal{E}|_L \longrightarrow 0.$$

Note that $\mathcal{E}|_L = \mathcal{E} \otimes \mathcal{O}_L$ is supported on a set of points. Taking cohomology gives a long exact sequence

$$\begin{array}{ccccccc} H^0(C, \mathcal{E}(-1)) & \rightarrow & H^0(C, \mathcal{E}) & \rightarrow & H^0(C, \mathcal{E}|_L) & \rightarrow & \\ \parallel & & & & & & \\ 0 & & & & & & \\ & & & & & & \\ H^1(C, \mathcal{E}(-1)) & \rightarrow & H^1(C, \mathcal{E}) & \rightarrow & H^1(C, \mathcal{E}|_L) & \rightarrow & \\ \parallel & & & & \parallel & & \\ 0 & & & & 0 & & \end{array}$$

This proves that $H^1(C, \mathcal{E}) = 0$.

It also proves $h^0(C, \mathcal{E}) = 2d$. Indeed, observe that $\mathcal{E}|_L = \mathcal{E} \otimes \mathcal{O}_L$ is of rank 2 supported on the set $C \cap L$ of d points, hence $h^0(C, \mathcal{E}|_L) = 2d$. $\square$

Theorem 3.6 is a special case of the following Beauville's corollary, where representations of hypersurfaces are studied via arithmetically Cohen-Macaulay (ACM) sheaves on $\mathbb{P}^n$. One of the advantages of our elementary proof is an explicit construction of representations from sheaves.

Theorem 3.7 (Corollary 2.4 in [1]). *Let X be an integral hypersurface of degree d in $\mathbb{P}^n(x_0, \dots, x_n)$ over a field k with $\text{char } k \neq 2$. Moreover, let E be an ACM vector bundle on X of rank 2 with determinant $\mathcal{O}_X(d+t)$. Then there exists a skew-symmetric matrix $A = (a_{ij}) \in \mathbb{M}_l$ with a_{ij} homogeneous polynomials in $x_0, \dots, x_n$ of degree $d_i + d_j - t$ and an exact sequence*

$$(8) \quad 0 \rightarrow \bigoplus_{i=1}^l \mathcal{O}_{\mathbb{P}^n}(t - d_i) \xrightarrow{A} \bigoplus_{i=1}^l \mathcal{O}_{\mathbb{P}^n}(d_i) \rightarrow E \rightarrow 0,$$

where X is defined by $\text{Pf } A = 0$.

If $H^0(X, E(-1)) = 0$ and $t = -1$, the entries of A are linear.

We conclude this section by Fujita's result [10, Example 6.4.16].

Remark 3.8. Suppose that C has genus $g \geq 2$. If $\mathcal{E}$ is a bundle on C such that $H^1(C, \mathcal{E}) = 0$, then $\mathcal{E}$ is ample. Being ample is equivalent to the condition that every quotient has strictly positive degree.

4. THE MODULI SPACE $M(2, 2(g-1))$

In this section we relate the set of pfaffian representations with the moduli space of semistable vector bundles. As described in Section 2, the idea arises from determinantal representations which can be parametrised by points on the Jacobian.

To any compact Riemann surface X one can associate the pair (JX, Θ) , the Jacobian and the Riemann theta function. The geometry of the pair is strongly related to the geometry of X . This gives an idea that higher rank vector bundles define a non-abelian analogue of the Jacobian called *moduli space* firstly due to the mathematicians of the Tata Institute. Much later physicists in Conformal Field Theory introduced pairs of moduli spaces and determinant line bundles on these moduli spaces. This has made a clear analogy with the Jacobian pair.

As before let C be a smooth plane curve of degree d and genus g . The existence and properties of

$$M_C(r, n),$$

the moduli space of semistable vector bundles on C of rank r and degree n , were established in [12], [15] and more modern treatment can be found in [14]. It is known that $M_C(r, n)$ is an irreducible, normal projective variety with an open subset $M_C^s(r, n)$ corresponding to stable bundles.

If C has genus $g \geq 2$ then $M_C^s(r, n)$ is not empty and its dimension is $r^2(g-1)+1$. The singular points of $M_C(r, n)$ are exactly $M_C(r, n) \setminus M_C^s(r, n)$.

If C is an elliptic curve then $M_C^s(r, n)$ is empty.

One can restrict the study to the moduli space

$$M_C(r, \mathcal{L})$$

of (semistable) rank r vector bundles on C with determinant $\mathcal{L}$. As described in [2], [5] it is a closed subvariety in $M_C(r, \deg \mathcal{L})$. Moreover, $M_C^s(r, \mathcal{L})$ is a closed subvariety in $M_C^s(r, \deg \mathcal{L})$. The determinant can be fixed since the moduli space $M_C(r, \deg \mathcal{L})$ is, up to a finite étale covering, the product of $M_C(2, \mathcal{L})$ with the Jacobian JC .

Drezed and Narasimhan [5] show that $\text{Pic}(M_C(r, \mathcal{L})) \cong \mathbb{Z}$ is generated by geometrically defined Cartier divisors $\Theta_{r, \mathcal{L}}$ in $M_C(r, \mathcal{L})$. For example, when $\deg \mathcal{L} = r(g-1)$ then

$$\chi(\mathcal{E}) = 0 \quad \text{for all } \mathcal{E} \in M_C(r, \mathcal{L})$$

and

$$\Theta_{r, \mathcal{L}} = \{ \mathcal{E} \in M_C(r, \mathcal{L}) : h^0(C, \mathcal{E}) \neq 0 \}$$

is naturally such a divisor.

Theorem 4.1. *Let C be a curve defined by a polynomial F of degree d in $\mathbb{P}^2$. There is a one to one correspondence between linear pfaffian representations of F (up to equivalence) and rank 2 bundles (up to isomorphism) on C in the open set*

$$M_C(2, \mathcal{O}_C(d-3)) \setminus \Theta_{2, \mathcal{O}_C(d-3)}.$$

Proof. From now on we consider the moduli space

$$M_C(2, \mathcal{O}_C(d-3))$$

of (semistable) rank 2 vector bundles on C with determinant $K_C \cong \mathcal{O}_C(d-3)$, which is a closed subvariety in $M_C(2, d(d-3)) = M_C(2, 2(g-1))$.

Recall from Theorem 3.6 the one to one correspondence between the linear pfaffian representations of C and rank 2 bundles on C with property (6) on page 8. For the sake of clearer notation, after tensoring by $\mathcal{O}_C(1)$, condition (6) can be rewritten into

$$(9) \quad \det \mathcal{E} \cong \mathcal{O}_C(d-3) \text{ and } H^0(C, \mathcal{E}) = 0.$$

Condition $h^0(C, \mathcal{E}) = 0$ implies that $\mathcal{E}$ is semistable. By the above considerations, bundles satisfying (9) can be parametrised by the points on

$$M_C(2, \mathcal{O}_C(d-3)) \setminus \Theta_{2, \mathcal{O}_C(d-3)}.$$

This is an open subset of $M_C(2, \mathcal{O}_C(d-3))$ since we cut out a Cartier divisor. $\square$

5. PFAFFIANS ARISING FROM DECOMPOSABLE VECTOR BUNDLES

In this section we find and explicitly describe linear pfaffian representations of C (up to equivalence) arising from decomposable vector bundles. We parametrise them by an open set in the Kummer variety. Since decomposable vector bundles are never stable [7, Ex V.2.8], this open set lies in the singular locus of $M_C(2, \mathcal{O}_C(d-3))$. The fact that the moduli space of rank 2 bundles with canonical determinant is singular along the Kummer variety can be found in [13].

Kummer variety $\mathcal{K}_C$ of C is by definition the quotient of the Jacobian JC by the involution $\mathcal{L} \mapsto \mathcal{L}^{-1} \otimes \mathcal{O}_C(d-3)$.




FIGURE 1. The darker areas represent the singular locus of M_C and decomposable Pfaffians respectively

Recall that by Theorem 3.6 and (9) we need to find all decomposable rank 2 bundles (up to isomorphism) with the property

$$\det \mathcal{E} \cong \mathcal{O}_C(d-3) \quad \text{and} \quad H^0(C, \mathcal{E}) = 0.$$

Write $\mathcal{E} = \mathcal{L} \oplus \mathcal{M}$. Then $\det \mathcal{E} \cong \mathcal{L} \otimes \mathcal{M} \cong \mathcal{O}_C(d-3)$ or equivalently

$$\mathcal{M} \cong \mathcal{L}^{-1} \otimes \mathcal{O}_C(d-3).$$

Observe that $\mathcal{E} \cong \mathcal{L} \oplus (\mathcal{L}^{-1} \otimes \mathcal{O}_C(d-3))$ has no sections if and only if $\mathcal{L}$ and $\mathcal{L}^{-1} \otimes \mathcal{O}_C(d-3)$ have no sections. These can be calculated by the Riemann-Roch formula and Serre duality

$$h^0(C, \mathcal{L}) - h^0(C, \mathcal{L}^{-1} \otimes \mathcal{O}_C(d-3)) = \deg \mathcal{L} + 1 - g,$$

since $\mathcal{O}_C(d-3)$ is exactly the line bundle of the canonical divisor K_C .

This proves that every decomposable rank 2 bundles with the property (9) is of the form

$$\mathcal{E} \cong \mathcal{L} \oplus (\mathcal{L}^{-1} \otimes \mathcal{O}_C(d-3)),$$

where $\mathcal{L}$ is a line bundle of degree $g-1 = \frac{1}{2}d(d-3)$ with no sections. These are exactly the *nonexceptional* line bundles defined in Section 2. We have seen that they correspond to the points on the Jacobian variety of C not on the exceptional subvariety W_{g-1} .

Conversely, by Section 2 a point on the Jacobian variety induces a determinantal representation M of C . Then

$$\begin{bmatrix} 0 & M \\ -M^t & 0 \end{bmatrix}$$

is a pfaffian representation of C with decomposable cokernel.

Thus we proved

Theorem 5.1. *There is a one to one correspondence between decomposable vector bundles in $M_C(2, \mathcal{O}_C(d-3)) \setminus \Theta_{2, \mathcal{O}_C(d-3)}$ and the open subset of Kummer variety*

$$(JC \setminus W_{g-1}) / \equiv,$$

where $\equiv$ is the involution $\mathcal{L} \mapsto \mathcal{L}^{-1} \otimes \mathcal{O}_C(d-3)$.

In the sequel explicitly construct the above correspondence from the sections of $\mathcal{E}(1)$. As above, $\mathcal{E} = \mathcal{L} \oplus (\mathcal{L}^{-1} \otimes \mathcal{O}_C(d-3))$ with $\mathcal{L}$ of degree $g-1$ and no sections. Denote $\mathcal{L} \otimes \mathcal{O}_C(1) = \mathcal{L}(1)$ by $\mathcal{F}$. Then

$$H^0(C, \mathcal{F}(-1)) = 0 \quad \text{and} \quad \deg \mathcal{F} = \frac{1}{2}d(d-1) = \deg(\mathcal{F}^{-1} \otimes \mathcal{O}_C(d-1)).$$

Let H be a section of $\mathcal{O}_C(1)$ and consider the exact sequence

$$0 \longrightarrow \mathcal{F}(-1) \longrightarrow \mathcal{F} \longrightarrow \mathcal{F}|_H \longrightarrow 0,$$

where $\mathcal{F}|_H$ is supported on a d points $C \cap H$. Applying cohomology gives a long exact sequence with dimensions

$$\begin{array}{ccccccc} H^0(C, \mathcal{F}(-1)) & \rightarrow & H^0(C, \mathcal{F}) & \rightarrow & H^0(C, \mathcal{F}|_H) & \rightarrow & H^1(C, \mathcal{F}(-1)) \rightarrow, \\ 0 & & & & d & & 0 \end{array}$$

since $h^0(C, \mathcal{F}(-1)) - h^1(C, \mathcal{F}(-1)) = 0$ by the Riemann-Roch theorem. This proves that $h^0(C, \mathcal{F}) = d$. The same way we show $h^0(C, \mathcal{F}^{-1} \otimes \mathcal{O}_C(d-1)) = d$. Let $\{f_1, \dots, f_d\}$ and $\{m_1, \dots, m_d\}$ be bases of the complete linear system of $\mathcal{F}$ and $\mathcal{F}^{-1} \otimes \mathcal{O}_C(d-1)$ respectively. Then

$$\{(f_1, 0), \dots, (f_d, 0), (0, m_1), \dots, (0, m_d)\}$$

form a basis for $H^0(C, \mathcal{E}(1))$. Obviously $(f_i, 0) \wedge (0, m_j) = f_i \otimes m_j$, whereas $(f_i, 0) \wedge (f_j, 0)$ and $(0, m_i) \wedge (0, m_j)$ equal 0. If we repeat the proof of Theorem 3.4, the map Ψ is of the form

$$\begin{bmatrix} 0 & f_i \otimes m_j \\ -f_j \otimes m_i & 0 \end{bmatrix}.$$

In coordinates it equals to a $2d \times 2d$ skew-symmetric matrix B with entries from the space of homogeneous polynomials of degree $d - 1$ and zero diagonal blocks. As before

$$A = \frac{1}{F^{d-2}} \tilde{B}$$

is a pfaffian representation of C . In particular,

$$A = \begin{bmatrix} 0 & M \\ -M^t & 0 \end{bmatrix},$$

where M is the determinantal representation of C corresponding to the nonexceptional line bundle $\mathcal{F}$.

6. CUBIC CURVE

In this section C will denote a curve defined by a smooth cubic polynomial F in $\mathbb{P}^2$.

Corollary 6.1. *On a cubic curve C all linear pfaffian representations can be parametrised by the points on the Kummer variety $\mathcal{K}_C - \{\text{one point}\}$.*

Proof. Recall that on an elliptic curve $K_C \cong \mathcal{O}_C$. Since $M_C^s(2, 0)$ is empty, there are no stable bundles on C . On the other hand, by [2, §4] the non-stable part of $M_C(2, \mathcal{O}_C)$ consists of decomposable vector bundles of the form $\mathcal{L} \oplus \mathcal{L}^{-1}$ for $\mathcal{L}$ in the Jacobian JC . But for $\mathcal{L} \in JC$ the following conditions are equivalent:

- $h^0(C, \mathcal{L} \oplus \mathcal{L}^{-1}) = 0$,
- $h^0(C, \mathcal{L}) = 0$,
- $\mathcal{L} \neq \mathcal{O}_C$.

Therefore

$$M_C(2, \mathcal{O}_C) \setminus \Theta_{2, \mathcal{O}_C} = \{\mathcal{L} \oplus \mathcal{L}^{-1}; \mathcal{L} \in JC\} \setminus \{\mathcal{O}_C \oplus \mathcal{O}_C\}.$$

□

Vinnikov [18] found an explicit one to one correspondence between the linear determinantal representations (up to equivalence) of C and the points on an affine piece of C :

Lemma 6.2 ([18]). *Every smooth cubic can be brought into the Weierstrass form*

$$F(x_0, x_1, x_2) = -x_1x_2^2 + x_0^3 + \alpha x_0x_1^2 + \beta x_1^3.$$

A complete set of determinantal representations of F is

$$x_0 \text{Id} + x_2 \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix} + x_1 \begin{pmatrix} \frac{s}{2} & l & \alpha + \frac{3}{4}s^2 \\ 0 & -s & -l \\ -1 & 0 & \frac{s}{2} \end{pmatrix},$$

where $l^2 = s^3 + \alpha s + \beta$. Note that the last equation is exactly the affine part $F(s, 1, l)$.

Recall that the Jacobian of a cubic curve C with $g = 1$ is the curve itself and $J - \{W_0\}$ is an affine piece of C . In particular, Corollary 6.1 implies that the complete set of pfaffian representations of F (put in the Weierstrass form) equals

$$\begin{bmatrix} 0 & M \\ -M^t & 0 \end{bmatrix},$$

where M are the determinantal representations in Lemma 6.2.

7. EXAMPLES OF HIGHER GENUS

We start by an example of a genus 3 curve.

Example 7.1. Any non hyperelliptic curve C of genus 3 is isomorphic to a plane quartic. In this case $M_C(2, \mathcal{O}_C(1)) \cong M_C(2, \mathcal{O}_C)$ embeds as a Coble quartic hypersurface in $\mathbb{P}^7$ and is singular along the Kummer variety $\mathcal{K}_C$. For references check [11], [9], [2].

For a given plane quartic Vanhaecke [17] gives explicit equations of the Coble quartic hypersurface. First he finds the equations of the Kummer variety which represents the singular locus of the moduli space, from here the Coble quartic is obtained by integration. For example, the moduli space of

$$C : x^4 - yz^3 - y^4 = 0$$

is the Coble hypersurface in $\mathbb{P}^7$ with equation

$$\begin{aligned} & z_0^4 - z_4^4 - 2z_1z_2^2z_3 - z_1^2z_3^2 - z_0z_3^3 - 4z_0z_2^2z_4 - 2z_0^2z_4^2 - 3z_0z_2z_4^2 + z_4^4 - 4z_0z_1z_2z_5 \\ & + z_3z_4z_5^2 + z_2z_5^3 - 4z_0z_1^2z_6 - 3z_0^2z_3z_6 - z_3z_4^2z_6 - 2z_2z_4z_5z_6 + z_1z_5^2z_6 - z_0z_6^3 - 2z_0^2z_1z_7 \\ & - z_3^2z_4z_7 + 2z_1z_4^2z_7 - z_2z_3z_5z_7 + z_0z_5^2z_7 - z_2^2z_6z_7 - z_1z_3z_6z_7 - 4z_0z_4z_6z_7 + z_1^2z_7^2. \end{aligned}$$

In the sequel we outline an algorithm for finding all pfaffian representations of the given C (up to equivalence) based on canonical forms of matrix pairs. This is a generalisation of Vinnikov's construction of determinant representations [18]. By Lancaster and Rodman [8, Theorem 5.1] every pfaffian representation of C can be put into the skew-symmetric canonical form

$$x \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ & & 0 & 0 & 0 & 1 & 0 & 0 \\ & & & 0 & 1 & 0 & 0 & 0 \\ & & & & 0 & 0 & 0 & 0 \\ & & & & & 0 & 0 & 0 \\ & & & & & & 0 & 0 \\ & & & & & & & 0 \end{pmatrix} + z \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ & & 0 & 0 & 1 & 0 & 0 & 0 \\ & & & 0 & 0 & 0 & 0 & 0 \\ & & & & 0 & 0 & 0 & 0 \\ & & & & & 0 & 0 & 0 \\ & & & & & & 0 & 0 \\ & & & & & & & 0 \end{pmatrix} + y \begin{pmatrix} 0 & c_{12} & c_{13} & c_{14} & c_{15} & c_{16} & c_{17} & c_{18} \\ & 0 & c_{23} & c_{24} & c_{25} & c_{26} & c_{27} & c_{28} \\ & & 0 & c_{34} & c_{35} & c_{36} & c_{37} & c_{38} \\ & & & 0 & c_{45} & c_{46} & c_{47} & c_{48} \\ & & & & 0 & c_{56} & c_{57} & c_{58} \\ & & & & & 0 & c_{67} & c_{68} \\ & & & & & & 0 & c_{78} \\ & & & & & & & 0 \end{pmatrix},$$

which we denote by A . Since $\text{Pf } A$ equals the equation of C we get

$$\begin{aligned}
c_{48} &= 1, \\
c_{15} &= -c_{18}^2 + 2c_{26}c_{28} + c_{28}c_{35} - c_{18}c_{36} - c_{36}^2 + 2c_{25}c_{38} + 2c_{18}c_{28}c_{38} - c_{28}c_{36}c_{38} - c_{26}c_{38}^2 - 2c_{35}c_{38}^2 - c_{18}c_{38}^3 \\
&\quad - c_{18}c_{45} - c_{36}c_{45} - c_{28}c_{38}c_{45} + c_{38}^3c_{45} - c_{45}^2 + c_{26}c_{46} - c_{28}^2c_{46} - c_{35}c_{46} + c_{18}c_{38}c_{46} + c_{36}c_{38}c_{46} + c_{28}c_{38}^2c_{46} \\
&\quad + 2c_{38}c_{45}c_{46} + c_{34}c_{56} + c_{24}c_{57} - c_{34}c_{38}c_{57} + c_{14}c_{58} - c_{28}c_{34}c_{58} - c_{24}c_{38}c_{58} + c_{34}c_{38}^2c_{58} + c_{23}c_{67} - c_{24}c_{38}c_{67} \\
&\quad + c_{13}c_{68} - c_{24}c_{28}c_{68} - c_{14}c_{38}c_{68} - c_{23}c_{38}c_{68} + c_{24}c_{38}^2c_{68} + c_{12}c_{78} - c_{14}c_{28}c_{78} - c_{13}c_{38}c_{78} + c_{14}c_{38}^2c_{78}, \\
c_{16} &= -c_{25} - c_{18}c_{28} + c_{28}c_{36} + c_{26}c_{38} + 2c_{35}c_{38} + c_{18}c_{38}^2 - c_{38}^2c_{45} - c_{18}c_{46} - c_{36}c_{46} - c_{28}c_{38}c_{46} \\
&\quad - 2c_{45}c_{46} + c_{34}c_{57} + c_{24}c_{58} - c_{34}c_{38}c_{58} + c_{24}c_{67} + c_{14}c_{68} + c_{23}c_{68} - c_{24}c_{38}c_{68} + c_{13}c_{78} - c_{14}c_{38}c_{78}, \\
c_{17} &= -c_{26} - c_{35} - c_{18}c_{38} + c_{38}c_{45} + c_{28}c_{46} + c_{34}c_{58} + c_{24}c_{68} + c_{14}c_{78}, \\
c_{37} &= -c_{28} - c_{46}, \\
c_{27} &= -c_{18} - c_{36} - c_{45}, \\
c_{47} &= -c_{38}.
\end{aligned}$$

There are 21 parameters c_{ij} left in the representation. We eliminate them by acting on A with an invertible constant matrix P via

$$P \cdot A \cdot P^t.$$

It is easy to check that P preserves the canonical form of A if and only if

$$P = \begin{bmatrix} Y^{-1} & Y^{-1}S \\ Y(T-S)^{-1} & Y(\text{Id} + (T-S)^{-1}S) \end{bmatrix},$$

where

$$Y = \begin{pmatrix} p & q & s & t \\ 0 & p & q & s \\ 0 & 0 & p & q \\ 0 & 0 & 0 & p \end{pmatrix}, \quad S = \begin{pmatrix} a & b & c & d \\ 0 & a & b & c \\ 0 & 0 & a & b \\ 0 & 0 & 0 & a \end{pmatrix}, \quad T = \begin{pmatrix} e & f & g & h \\ 0 & e & f & g \\ 0 & 0 & e & f \\ 0 & 0 & 0 & e \end{pmatrix}.$$

This enables us to reduce to 12 parameters c_{ij}

$$\begin{bmatrix} 0 & c_{12} & c_{13} & 0 & -c_{36}^2 - c_{36}c_{45} - c_{45}^2 + c_{26}c_{46} - c_{35}c_{46} + c_{23}c_{67} & -c_{25} - c_{36}c_{46} - 2c_{45}c_{46} & -c_{26} - c_{35} & 0 \\ & 0 & c_{23} & 0 & c_{25} & c_{26} & -c_{36} - c_{45} & 0 \\ & & 0 & 0 & c_{35} & c_{36} & -c_{46} & 0 \\ & & & 0 & c_{45} & c_{46} & 0 & 1 \\ & & & & 0 & c_{56} & c_{57} & 0 \\ & & & & & 0 & c_{67} & 0 \\ & & & & & & 0 & 0 \\ & & & & & & & 0 \end{bmatrix}$$

Another action by P which preserves the 0 and 1 elements in the above matrix yields

$$\begin{aligned}
 c_{12} &\mapsto \frac{c_{12}+a(-2c_{25}-2c_{36}c_{46}-3c_{45}c_{46}+ac_{56}+ac_{46}c_{67})}{p^2} \\
 c_{13} &\mapsto \frac{c_{13}+a(-c_{26}-2c_{35}-c_{46}^2+ac_{57})}{p^2} \\
 c_{23} &\mapsto \frac{c_{23}+a(-2c_{36}-c_{45}+ac_{67})}{p^2} \\
 c_{25} &\mapsto \frac{c_{12}-ac_{25}-c_{23}c_{46}-2ac_{45}c_{46}-ec_{25}+aec_{56}}{a-e} \\
 c_{26} &\mapsto c_{26} \\
 c_{35} &\mapsto \frac{c_{13}-ec_{35}-a(c_{26}+c_{35}+c_{46}^2-ec_{57})}{a-e} \\
 c_{36} &\mapsto \frac{c_{23}-ec_{36}-a(c_{36}+c_{45}-ec_{67})}{a-e} \\
 c_{45} &\mapsto c_{45} \\
 c_{46} &\mapsto c_{46} \\
 c_{56} &\mapsto \frac{(c_{12}-c_{23}c_{46}+e(-2c_{25}-2c_{45}c_{46}+ec_{56}))p^2}{(a-e)^2} \\
 c_{57} &\mapsto \frac{(c_{13}+e(-c_{26}-2c_{35}-c_{46}^2+ec_{57}))p^2}{(a-e)^2} \\
 c_{67} &\mapsto \frac{(c_{23}+e(-2c_{36}-c_{45}+ec_{67}))p^2}{(a-e)^2}
 \end{aligned}$$

Observe that $c_{13}c_{57} - c_{35}(c_{35} + c_{26} + c_{46}^2)$, $c_{23}c_{67} - c_{36}(c_{36} + c_{45})$ are invariant under this action.

We are left with 12 parameters c_{ij} modulo $P(a, e, p)$ action. The condition $\text{Pf } A = x^4 - yz^3 - y^4$ gives three additional equations

$$\begin{aligned}
 &2c_{36}^2c_{45}+2c_{36}c_{45}^2+c_{45}^3+2c_{35}c_{45}c_{46}+c_{25}(c_{26}+2c_{35}+c_{46}^2)+c_{23}c_{46}c_{57} = \\
 &2c_{26}c_{45}c_{46}+c_{13}c_{56}+c_{12}c_{57}+2c_{23}c_{45}c_{67}, \\
 &c_{36}^4+2c_{36}^3c_{45}+c_{25}^2c_{46}+2c_{25}c_{45}c_{46}^2+c_{26}^2(-c_{35}+c_{46}^2)+c_{13}c_{45}c_{56}+c_{23}c_{25}c_{57}+2c_{23}c_{45}c_{46}c_{57}+c_{12}c_{35}c_{67}+ \\
 &2c_{23}^2c_{67}^2+2c_{36}^2(c_{45}^2-c_{23}c_{67})+c_{36}(c_{45}^3+c_{25}c_{46}^2+c_{13}c_{56}-c_{12}c_{57}+c_{23}c_{46}c_{57}-2c_{45}(c_{35}c_{46}+c_{23}c_{67})) = \\
 &1+c_{25}c_{35}c_{45}+2c_{35}c_{45}^2c_{46}+2c_{35}c_{35}c_{56}+c_{12}c_{46}c_{56}+c_{13}c_{25}c_{67}+c_{23}c_{45}^2c_{67}+c_{23}c_{35}c_{46}c_{67}+ \\
 &c_{26}(c_{35}^2-c_{25}c_{36}+2c_{36}^2c_{46}+2c_{36}c_{45}c_{46}+c_{45}^2c_{46}+c_{35}c_{46}^2+c_{23}c_{56}-c_{13}c_{57}-2c_{23}c_{46}c_{67}), \\
 &c_{26}^2+c_{26}c_{35}+c_{35}^2+2c_{25}c_{36}+c_{25}c_{45}+2c_{36}^2c_{46}+4c_{36}c_{45}c_{46}+3c_{45}^2c_{46}+c_{35}c_{46}^2 = \\
 &c_{26}c_{46}^2+c_{13}c_{57}+c_{12}c_{67}+c_{23}(c_{56}+c_{46}c_{67}).
 \end{aligned}$$

We proved that all pfaffian representations of the given quartic curve form a $6 = 12 - 3(\text{parameters } a, e, p) - 3(\text{relations among } c_{ij})$ dimensional affine variety.

Let C be a generic plane curve of genus $g \geq 3$. By [13] the moduli space $M_C(2, \mathcal{O}_C(d-3))$ of rank 2 bundles with canonical determinant embeds into $|2\Theta|$. Its singular locus is isomorphic to the Kummer variety. Moreover, the embedding restricts to the Kummer map on the singular locus.

There are not many results in the literature explicitly describing the above moduli spaces for curves of genus $g > 3$. Finding all pfaffian representations of a given curve C would provide a description of the open set $M_C(2, \mathcal{O}_C(d-3)) \setminus \Theta_{2, \mathcal{O}_C(d-3)}$.

Vanhaecke's method could be used for curves of higher genus. Start by finding all determinantal representations of C (which are smaller than Pfaffians) using

Vinnikov's method of canonical forms [19]. Determinantal representations induce pfaffian representations of the form

$$\begin{bmatrix} 0 & \diamond \\ -\diamond^t & 0 \end{bmatrix},$$

which by Section 5 correspond to decomposable bundles. These define the singular locus of the moduli space. The equations of the moduli space could be then found by integration.

REFERENCES

- [1] A. Beauville. *Determinantal Hypersurfaces*, Michigan Math. J. **48** (2000), 39–63.
- [2] A. Beauville. *Vector bundles on curves and generalized theta functions: recent results and open problems*, Complex Algebraic Geometry, MSRI Publications **28** (1995), 17–33.
- [3] R. J. Cook and A. D. Thomas. *Line bundles and homogeneous matrices*, Quart. J. Math. Oxford Ser. (2) **30** (1979), 423–429.
- [4] I. Dolgachev. *Topics in classical algebraic geometry*, Lecture Notes <http://www.math.lsa.umich.edu/~idolga/lecturenotes.html>.
- [5] J.M. Drezet and M.S. Narasimhan. *Groupe de Picard des varietes de modules de fibres semi-stables sur les courbes algebriques*, Invent. Math. **97** (1989), 53–94.
- [6] W. Fulton and P. Pragacz. *Shubert varieties and degeneracy loci*, Lecture notes in Mathematics **1689**, Springer-Verlag, 1998.
- [7] R. Hartshorne. *Algebraic Geometry*, Graduate Texts in Mathematics **52**, Springer-Verlag, 1977.
- [8] P. Lancaster and L. Rodman. *Canonical forms for symmetric / skew-symmetric real matrix pairs under strict equivalence and congruence*, Linear Algebra Appl. **406**, (2005), 1–76.
- [9] Y. Laszlo. *A propos de l'espace des modules des fibres de rang 2 sur une courbe*, Math. Annalen **299**, (1994), 597–608.
- [10] R. Lazarsfeld. *Positivity in Algebraic Geometry II*, A Series of Modern Surveys in Mathematics **49**, Springer-Verlag, 2000.
- [11] M.S. Narasimhan and S. Ramanan. *2Θ linear systems on Abelian varieties*, Vector bundles on algebraic varieties, Oxford University Press (1987), 415–427.
- [12] P. E. Newstead. *Introduction to moduli problems and orbit spaces*, Tata Institute of Fundamental Research, Bombay, Springer-Verlag, 1978.
- [13] W. M. Oxbury, C. Pauly and E. Previato. *Subvarieties of SU and 2Θ-divisors in the Jacobian*, Transactions of the AMS Vol 350, **9**, (1998).
- [14] J. Le Potier. *Lectures on Vector Bundles*, Cambridge studies in advanced mathematics **54**, Cambridge University Press, 1997.
- [15] C. S. Seshadri. *Fibres vectoriels sur les courbes algebriques*, Asterisque **96**, 1982.
- [16] I. R. Shafarevich. *Basic Algebraic Geometry 1*, Springer-Verlag, 1994.
- [17] P. Vanhaecke. *Integrable systems and moduli spaces of rank two vector bundles on a non-hyperelliptic genus 3 curve*, Ann. Inst. Fourier 55, **6**, (2005), 1789–1802.
- [18] V. Vinnikov. *Complete description of determinantal representations of smooth irreducible curves*, Lin. Alg. Appl., **125** (1989), 103–140.
- [19] V. Vinnikov. *Determinantal representations of real cubics and canonical forms of corresponding triples of matrices*, Mathematical Theory of Networks and Systems, Lecture Notes in Control and Inform. Sci. 58, 1984.
- [20] M. Marcus and R. Westwick. *Linear maps on skew-symmetric matrices: The invariance of elementary symmetric functions*, Pacific J. Math. (3) **14** (1960).

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF LJUBLJANA, JADRANSKA 19, 1000 LJUBLJANA, SLOVENIA e-mail: anita.buckley@fmf.uni-lj.si